SAMPLE APPROVED PROTOCOLS

ADMINISTRATION OF ASPIRIN FOR CHEST PAIN BELIEVED TO BE OF CARDIAC ORIGIN

PROVIDER NAME:			: PROVIDER NO. 60
havii	ng chest	pain be	e used by properly trained and licensed EMTs for the treatment of patients who are lieved to be of cardiac origin. EMTs-basic must obtain a direct voice order from the ician in all cases prior to administration of aspirin.
I.	Aspir	in	
	A.		raindications
		1.	Inability to swallow
		2.	Allergy to Aspirin
		3.	Patient has already taken the maximum recommended dose prior to EMT arrival.
II.	CHEST PAIN BELIEVED TO BE OF CARDIAC ORIGIN - Patient presents with complaint or		
	chest	-	
	A.		rm initial assessment
	B.		n patient history and perform physical exam for cardiac patient
		1.	History of chest pain? Heart disease?
		2.	Onset of chest pain and progression since onset?
	C	3.	Interventions (previous medications taken, aspirin, nitroglycerin)
	C. D.	Assess baseline vital signs and SAMPLE history Administer oxygen (if not already done during initial assessment) - consider Advanced	
	D.	Life Support	
	E.	Monitor patient using monitoring electrodes per local protocol	
	F.	Contact medical control for authorization to administer between 160-325 mg. of aspir	
	orally		
		1.	Obtain voice authorization for administration, including dosage; repeat order back
			to physician
		2.	Verify medication and that medication has not expired
		3.	Describe procedure to patient and obtain consent
		4.	Administer medication
		5.	Record actions and reassess patient
			or aspirin disapproved - transport, continue to monitor patient and maintain
cont			al control.
	G.		port and continue to monitor and reassess patient
	H.	Kepo	rt any changes to hospital
Approved by:		: _	Medical Director (Print)
			Medical Director (Signature)

Date